

The Determination of Internally Deposited Radioactive Isotopes  
in the  
Marshallese People  
by  
Excretion Analysis<sup>#</sup>

Kent T. Woodward, Ariel G. Schrod<sup>##</sup>t, James E. Anderson,  
Harry A. Claypool, and James B. Hartgering

Division of Nuclear Medicine  
Walter Reed Army Institute of Research  
Washington 12, D. C.

**BEST COPY AVAILABLE**

RG
Collection 1320
Box 9
Folder 43

<sup>#</sup> Work done under the auspices of The Surgeon General, United States Army, and in conjunction with the Division of Biology and Medicine, Atomic Energy Commission

<sup>##</sup> Present Address: Nuclear-Chicago, Chicago, Illinois

## INTRODUCTION.

Following the detonation of a thermonuclear device at the Pacific Test Site on 1 March 1954, 239 Marshallese people were exposed to significant levels of gamma radiation from fallout. Estimated total exposures ranged from 175r on Rongelap to 14r on Utirik (1).

These populations were evacuated to Kwajalein for decontamination and care. During the two days of fallout exposure before evacuation was completed, the Marshallese also received some radioactive materials internally by ingestion and inhalation. Estimates of the internal body burden from fallout were obtained from the analysis (1) of urine samples collected soon after exposure.

These data indicated that the acute hazard from internally deposited fission fragments was quite small as compared to the whole body gamma radiation exposure. Although the radioactivity levels in the urine were low, the activity was sufficient to obtain reasonable precision and to warrant additional long term studies of the activity levels and excretion patterns of this rather large and well isolated population.

The people from Alinginae and Utirik were returned to their home islands in June 1954. Radiation intensities on Rongelap, however, precluded an early return to this atoll and the Rongelap people lived on Majuro from June 1954 until July 1957.

Basic data on the food crops of the Marshallese indicated that after the resettlement on the contaminated atolls, intake of strontium<sup>90</sup> would be increased considerably, and that cesium<sup>137</sup>, zinc<sup>65</sup>, and cobalt<sup>60</sup> were dietary constituents of island and ocean foodstuffs, and also would be assimilated (2). The expected increases in the trace amounts of radionuclides in the food supply of a large population would afford an opportunity to investigate the rate of equilibration and the discrimination factors operating between food supply and man. Urinary

DOE ARCHIVES

excretion levels of cesium<sup>137</sup> and strontium<sup>90</sup> were measured from March 1954 through March 1958. Zinc<sup>65</sup> levels were first measured in 1958 samples.

#### MATERIALS AND METHODS.

One hundred and forty one individual urine samples collected from 24 March 1954 through 7 September 1954 were obtained by the Health and Safety Laboratory, AEC. Urine volumes were small (about 350 ml) and it was necessary to pool samples. This was done according to the age of the subjects and 19 samples of pooled urine were assayed. A 57 liter pooled urine sample from Rongelap was collected and assayed in 1956 (3). Three pooled samples and seven individual samples were assayed in 1957. Thirty individual urine samples were assayed in 1958.

In samples collected in 1954 and 1957 cesium<sup>137</sup> was scavenged by nickel ferrocyanide (urine made strongly alkaline) and counted in a crystal well counter. A twenty channel gamma-ray spectrum was determined for each sample and the cesium<sup>137</sup> photo spectrum count rate used. The 1958 samples were assayed directly for Cs<sup>137</sup>, Zn<sup>65</sup>, and K<sup>40</sup> in 2.5 liter plastic containers placed on an 8 x 4 inch (TH activated) sodium iodide crystal. The activity for each radio-isotope was determined by gamma-ray spectral analysis. Sample activities were compared with known radioactive standards ( $\pm 5$  percent) counted in the same geometry.

Strontium<sup>90</sup> was precipitated from urine as the carbonate. Yttrium<sup>90</sup> was separated and identified by its half-life using thin walled gas flow counters.

Urine samples were corrected for radio-active decay to the time of collection.

There is some uncertainty as to the completeness and the duration of time over which samples were collected and therefore twenty-four hour urine volumes are not accurately known. Potassium<sup>40</sup> excretion, using 360 d/m or 2 gm K/day indicates an average daily volume of about 1180 ml ( $\pm 56$  percent). It was convenient to use one liter as an average 24 hour urine volume and to express radio-assays in micromicrocuries per liter.

## RESULTS AND DISCUSSION.

### Cesium<sup>137</sup> Excretion Levels and Body Burden

The urinary excretion levels of cesium<sup>137</sup> for the years 1954, 1957, and 1958 are shown in Tables 1 - 4. On 24 - 25 March, 1954, the mean excretion level of cesium<sup>137</sup> for all age groups was 405 uuc per liter. With an excretion rate of 0.46 percent (4) of cesium<sup>137</sup> body burden per 24 hours, the mean body burden from fallout 24 - 25 days after exposure was  $405/4.6 \times 10^{-3} \times 10^6$  or 88 muc ( $\pm 54$  percent). This value is about 20 times the average body burden reported by E. C. Anderson, et al (5, 6) for people measured during 1956 - 1957 in the United States. The cesium<sup>137</sup> urinary excretion levels for the six months following exposure can be expressed as an exponential function, and a best line of fit drawn through the data resulted in a half time for elimination of about 110 days (Fig 1). A biological half time of about 140 days has been observed on volunteers who ingested one microcurie of radio-cesium (4).

From the 1957 Cs<sup>137</sup> excretion levels (Table 2) the Rongelap group exposed to fallout was estimated to have an average burden of about 7 muc, whereas the Rongelap control group was about 2 muc. Body burden in either group in 1957 is comparable to levels measured in the U. S. population (6). With a half time for elimination of the order of 150 days, the body burden of the exposed Rongelap group should have decreased from the March 1954 level to 7 muc in about 550 days, or late in 1955. A body burden of 7 muc for this group in March 1957 could then indicate a continuing exposure to Cs<sup>137</sup> during 1956 of the order of 32 micro-microcuries per day from stratospheric-tropospheric fallout while residing on Majuro.

Since the Utirik group was returned to their atoll in 1954, the mean body burden in 1957 was elevated to an estimated 337 muc, some 48 times the Cs<sup>137</sup> burden of the exposed Rongelap people who resided on Majuro. This long residency time on Utirik atoll after fallout contamination, as compared to the excretion

rate of  $\text{Cs}^{137}$  should have resulted in an equilibrated  $\text{Cs}^{137}$  burden, with an estimated daily intake of about 1560 uuc of  $\text{Cs}^{137}$ . Unfortunately no systematic survey of foodstuffs grown on these atolls has been reported. Data available, however show that coconut grown on Rongelap contained about 9 uuc  $\text{Cs}^{137}$  per gram, and arrowroot (Utirik) contained about 8 uuc  $\text{Cs}^{137}$  per gram. The daily intake of several hundred grams of either staple would be sufficient to account for the 1957 excretion level in the Utirik group.

The Rongelap groups had been resettled for about nine months at the time of the March 1958 medical survey, and urinary excretion levels of  $\text{Cs}^{137}$  had increased about one hundred fold over 1957 levels. Mean body burden for the two groups at this time was 0.9 uc ( $\pm 27$  percent) and 1.2 uc ( $\pm 47$  percent) (Tables 3 and 4). Cesium<sup>137</sup> body burden may have equilibrated by late 1958 and predicted burdens were about 1.3 and 1.6 uc respectively.

#### Strontium<sup>90</sup> Excretion Levels and Body Burden

Urinary excretion levels of strontium<sup>90</sup> are presented in Tables 1, 2, and 3. The strontium<sup>90</sup> excretion level in 1956 was 0.5 uuc/liter as determined in a pooled sample of 57 liters. Figure 2 shows the excretion of  $\text{Sr}^{90}$  for the three years following fallout exposure. Although there is considerable variation in the data for the various age groups at early times, mean values for all groups plotted suggest that the excretion pattern can be expressed conveniently as the sum of two exponential terms. The larger portion of  $\text{Sr}^{90}$  was excreted with a half time of about 40 days, and a small fraction, 20 percent, was excreted with a half time of about 500 days. This is similar to Cowan's (7) urinary excretion study of an accident case involving inhaled  $\text{Sr}^{90}$ .

As was noted in the March 1958  $\text{Cs}^{137}$  levels, the excretion levels of  $\text{Sr}^{90}$  were also increased to 3.5-4.0/0.2, or about 20 fold. Since  $\text{Cs}^{137}$  levels increased 4300 - 5300/34, or about 140 fold, the ratio is about seven in favor of cesium<sup>137</sup>. With the increases in urinary  $\text{Sr}^{90}$  excretion levels in 1958, it was

pertinent to estimate body burden, burden expected at equilibrium, and daily intake of Sr<sup>90</sup> from these excretion levels.

The metabolic behavior of strontium as outlined in Supplement #6 of the British Journal of Radiology was used to estimate body burden, etc. from urinary excretion levels of strontium<sup>90</sup> (Appendix). The fraction of strontium absorbed from the gastro-intestinal tract is 0.6 and the biological excretion rate from the total body is 190 days. Of the absorbed fraction, 0.25/0.60, about 42 percent is deposited in bone and the biological half-life is 4000 days. Assuming that the absorbed fraction is excreted entirely in urine, the mean body burden of the exposed Rongelap group in March 1958 was 2 muc ( $\pm$  52 percent). This is about nine percent of the expected equilibrium value of 23 muc. The estimated burden of strontium<sup>90</sup> for March 1958 is probably too low and compares with levels measured in stillborn children in the U. S. several years ago (8). The daily intake of strontium<sup>90</sup> is estimated to be about 15 micromicrocuries or 15 Sunshine Units (assuming a daily calcium intake of one gram).

Dunning (2) reported that the average concentration of strontium<sup>90</sup> in the Marshallese food supply could be about 360 Sunshine Units, but this would reduce to well under 100 Sunshine Units if the consumption of high Sr<sup>90</sup> content foods were eliminated. With the elimination of pandanus and land crabs the diet used by Dunning indicated that the intake of strontium<sup>90</sup> would be 17 Sunshine Units per day. This compares favorably with the estimated intake of about 15 micromicrocuries from excretion analysis.

#### Zinc<sup>65</sup> Excretion Levels and Body Burden

In early 1957 Miller (9) detected Zn<sup>65</sup> in selected residents of Rongelap and Utirik by whole body gamma-ray spectrometry. Body burden ranged from 29.5 to 73.0 muc for the Rongelap residents, and 482 and 229 muc was detected in two subjects from Utirik. The Rongelap subjects were residing on Majuro at this time.

Miller obtained an effective half-time of 110 days for the elimination of  $Zn^{65}$ , and for the two subjects from Utirik the urinary to fecal excretion ratio was 1/9.

Assuming the excretion to be entirely exponential and 10 percent of the body burden of  $Zn^{65}$  excreted in urine, the March 1958 urinary excretion levels of 174 and 342 micromicrocuries indicate body burden, equilibrium body burden, and daily intake as follows:

	RONGELAP	
	1954 Exposed Group	Control Group (Unexposed 1954)
Body Burden (March 1958):	280 muc ( $\pm 49\%$ )	540 muc ( $\pm 90\%$ )
Equilibrated Body Burden:	330 muc	650 muc
Daily Intake:	2100 uuc/day #	4100 uuc/day #
Percent Equilibration:	85.0 percent	83.0 percent

The mean body burden estimated from 1958 excretion analysis for all Rongelap subjects showed a ten-fold increase over the 1957 whole body measurements. This increase correlates with the return of these people to Rongelap atoll from Majuro. Also the 1958 Rongelap  $Zn^{65}$  burdens are comparable with the Utirik subjects in 1957, and the Utirik subjects would have been in equilibrium in 1957 (half time of 110 days for the elimination of zinc<sup>65</sup>).

The estimated intake of zinc<sup>65</sup> (2000 to 4000 uuc per day) can not be accounted for by  $Zn^{65}$  activity levels reported in foodstuffs. Although this radio-nuclide reportedly accounts for a large fraction of the total activity in fish, this amounts to only about six uuc per pound of muscle up to 75 uuc per pound of whole fish (2) or at most four percent of the estimated intake.

#### CONCLUSIONS.

Since resettlement of the Marshallese people on Rongelap atoll in July 1957, the urinary excretion level of cesium<sup>137</sup> has increased about 140 fold and about

# Assuming 100 percent absorption from the GI Tract

20 fold for strontium<sup>90</sup>. Zinc<sup>65</sup> was readily detected in samples from the March 1958 medical survey.

The estimated mean body burden at equilibrium for cesium<sup>137</sup> is about 1.5 microcuries or about 1/6 of the tolerance recommended by the International Commission for Radiological Protection for non-industrial populations. For strontium<sup>90</sup> the mean body burden of the exposed Rongelap group in March 1958 was estimated to be two millimicrocuries. This is about nine percent of the expected equilibrium value of 23 millimicrocuries. The equilibrated strontium<sup>90</sup> burden is about 1/5 of tolerance. The estimated mean body burden of zinc<sup>65</sup> for Rongelap subjects in March 1958 is about 85 percent of the equilibration value of 0.6 microcuries and the equilibration value is 1/70 of tolerance.



## BIBLIOGRAPHY

1. Cronkite, E. P., V. P. Bond, C. L. Dunham, etc.  
Some Effects of Ionizing Radiation on Human Beings, Atomic Energy Commission, July (1956); WT 923 (1954)
2. Dunning, G. M.  
Radioactive Contamination of Certain Areas in the Pacific Ocean from Nuclear Tests, Atomic Energy Commission, August (1957)
3. Conard, R. A., B. Cannon, C. E. Huggins, J. B. Richards, and A. Lowery.  
Medical Survey of Marshallese Two Years Post Exposure to Fallout Radiation, Brookhaven National Laboratory, BNL-412 (T-80), September (1956)
4. Woodward, K. T., C. R. Richmond, and W. H. Langham .  
Measurements of Retention and Excretion of Radioisotopes of the Alkali Metals by Mice and Rats, Using an Annular Liquid Scintillation Counter, Proc. Health Phys. Soc. 79 (1956)
5. Anderson, E. C., R. L. Schuch, W.R. Fisher, and W. H. Langham.  
Radioactivity of People and Foods, Science, 125:1273 (1957)
6. Anderson, E. C.  
Radioactivity of People and Milk: 1957, Science, 128:882 (1958)
7. Cowan, F. P., L. B. Farabee, and R. A. Love.  
Health Physics and Medical Aspects of a Strontium<sup>90</sup> Inhalation Accident. Am. J. Roentgenol., Rad. Ther., and Nuc. Med, 67:805 (1952)
8. Libby, W. F.  
Current Research Findings on Radioactive Fallout. Proc. Nat. Acad. Sci. (Wash) 42, 945 (1956)
9. Miller, C. E.  
Measurements on Some Residents of the Marshall Island, Argonne National Laboratory, Report No. ANL 5755, pg 56 (1957)

TABLE 1

EXCRETION LEVELS OF URINARY CESIUM<sup>137</sup> AND STRONTIUM<sup>90</sup>  
(uuc per liter) IN THE MARSHALLESE AT VARIOUS TIMES AFTER EXPOSURE IN 1954

DATE OF COLLECTION	AGE GROUPS										MEAN ±S.D.	
	< 5 yrs		5-16 yrs		16-24 yrs		24-40 yrs		> 40 yrs		Cs <sup>137</sup>	Sr <sup>90</sup>
	Cs <sup>137</sup>	Sr <sup>90</sup>	Cs <sup>137</sup>	Sr <sup>90</sup>	Cs <sup>137</sup>	Sr <sup>90</sup>	Cs <sup>137</sup>	Sr <sup>90</sup>	Cs <sup>137</sup>	Sr <sup>90</sup>		
24 March 1954	-	-	889	11.0	294	5.4	372	3.9	258	-	405±218	7.1±2.4
25 March 1954	-	-	-	-	-	-	268	7.7	352	7.7		
17 April 1954	794	16.4	780	-	431	1.7	311	1.2	323	2.3	528±214	5.4±6.5
14 or 31 May 1954	-	-	255	13.4	427	4.2	434	0.9	543	5.5	415±100	6.0±4.6
6 or 7 Sep 1954	-	-	118	2.0	281	1.9	86	0.5	141	0.5	157±73	1.2±0.6

DOE ARCHIVES

TABLE 2

EXCRETION LEVELS OF URINARY CESIUM<sup>137</sup> AND STRONTIUM<sup>90</sup>  
IN THE MARSHALLESE DURING MARCH 1957

SOURCE	MEAN SAMPLE VOLUME	ACTIVITY (mc/liter)	
		CESIUM <sup>137</sup>	STRONTIUM <sup>90</sup>
Exposed - Rongelap	4,100 ml	34.	0.2
Controls - Rongelap	3,664 ml	8.	< 0.2
Exposed - Utirik	2,875 ml	1535.	0.2
	TOTAL SAMPLE VOLUME		
BILLET #9	5,400 ml	62.	0.5
IROJI #26	10,200 ml	168.	0.6
JOHN #40	2,700 ml	128.	-
TIMA #79	5,400 ml	103.	< 0.2
TOTAK #82	8,800 ml	120.	< 0.2
ALEK #2123	2,700 ml	3,759.	-
LEBAN #2125	5,400 ml	1,698	< 0.2

DOE ARCHIVES

TABLE 3  
EXCRETION LEVELS OF URINARY CESIUM<sup>137</sup>, POTASSIUM<sup>40</sup>, ZINC<sup>65</sup>, and STRONTIUM<sup>90</sup> DURING MARCH 1958

1954-EXPOSED GROUP  
RONGELAP

SUBJECT	CASE NO	SEX	AGE (1958)	URINE VOL. (ml)	ACTIVITY				
					CESIUM <sup>137</sup> uuc/l	POTASSIUM <sup>40</sup> gm K/l	Cs/K uuc/gm	ZINC <sup>65</sup> uuc/l	STRONTIUM <sup>90</sup> uuc/l
BELLA	7	M	41	2680	2181	1.0	2203	162	1.6
BILLIET	9	M	27	5700	1233	0.7	1665	100	3.8
AIMIRA	12	F	23	6745	2924	1.3	2232	264	1.5
ETRI	22	F	21	5525	5917	2.5	2357	345	6.0
IROJI	26	M	16	5915	4330	1.6	2706	223	2.1
JANWOR	31	M	36	2580	3393	2.3	1488	238	1.2
JIMAKO	39	F	19	130	13130	--	--	155	NDA
JOHN	40	M	34	1740	2275	0.9	2615	148	6.1
JOJIA	41	M	48	2690	2245	--	--	107	5.3
MONNA	66	F	34	2665	2413	1.4	1664	22	3.1
NIKTIMOS	73	M	22	4125	5584	--	--	147	5.7
NORIO	76	M	13	2665	11708	0.3	45031	237	2.8
TIMA	79	M	49	1015	3717	2.1	1796	121	2.0

DOE ARCHIVES


TABLE 4

EXCRETION LEVELS OF URINARY CESIUM<sup>137</sup>, POTASSIUM<sup>40</sup>, ZINC<sup>65</sup>, and STRONTIUM<sup>90</sup> DURING MARCH 1958CONTROL GROUP  
(UNEXPOSED-1954)  
RONGELAP

SUBJECT	CASE NO.	SEX	AGE (1958)	URINE VOL. (ml)	CESIUM <sup>137</sup>	POTASSIUM <sup>40</sup>	Cs/K	ZINC <sup>65</sup>	STRONTIUM <sup>90</sup>
					uuc/1	gm K/1	uuc/gm	uuc/1	uuc/1
TOLMIA	818	M	7	1880	7674	0.3	24755	99.	6.4
REKO	825	F	16	400	9928	2.8	3546	337.	10.2
KEJAI	830	M	20	4275	5165	1.9	2662	553.	6.7
LOLE	831	M	18	1430	7342	1.4	5063	306	2.7
BWIO	836	M	24	585	7028	3.8	1835	88.	2.5
SAMSEN	838	M	26	10515	1867	1.1	1652	324.	3.5
ARTICLE	840	M	31	2355	3393	1.4	2458	1262.	4.1
LATA	843	F	33	6490	2068	0.7	3041	75.	1.7
NAMON	849	M	39	2640	3880	3.3	1158	948.	5.0
MONEAN	855	M	60	2655	3176	1.2	2669	120.	2.5
JONNI	865	F	25	2125	4624	1.7	2688	319.	4.1
BATOL	872	M	14	5275	7736	1.9	3947	195.	1.3
ANTY	874	M	10	4650	6141	0.9	6533	131.	3.2
HANAKO	876	F	20	2155	2909	0.3	9091	163.	3.3
ATARIK	877	M	20	5245	3311	1.5	2164	233.	2.0
JEKOB	883	M	46	2615	3386	2.1	1628	271.	2.7
KETO	887	M	13	2630	11733	2.2	5357	398	4.7

DOE ARCHIVES

# CESIUM <sup>137</sup> EXCRETION (μμc/LITER)



DOE ARCHIVES

Photo No. C484-1

Figure 1. Excretion Levels of Urinary Cesium<sup>137</sup> at Various Times After Exposure

DOE ARCHIVES

STRONTIUM<sup>90</sup> EXCRETION ( $\mu\mu\text{c/LITER}$ )






Photo No. 9648-2

Figure 2. Excretion Levels of Urinary Strontium<sup>90</sup> at Various Times After Exposure

DOE ARCHIVES

APPENDIX

I. In the case of strontium elimination, the following assumptions were made:

a. The population was returned to the contaminated atoll at time  $t = 0$ , with a zero strontium body burden.

b. The population absorbs a daily increment of  $x$  uc, and  $x$  is considered to be a constant independent of time.

c. The amount of strontium excreted in the urine each day is given by  $P(t) = \frac{1}{k}E(t)$ , where  $E(t)$  is the total excreted by all routes each day, and  $k$  is a constant independent of time.

d. The body is considered to be a two compartment system, A and B, where  $A + B = 1$ . The excretion rates for each compartment are  $a$  and  $b$  days<sup>-1</sup> respectively. The portion of  $E(t)$  excreted from each compartment is proportional to the burden remaining in that compartment. For cesium and zinc elimination similar assumptions are made, except that only one compartment is assumed.

e. Now:

$s(t)$  is total strontium body burden at time  $t$  in uc

$s_1(t)$  and  $s_2(t)$  are the portions in each compartment

and  $s(t) = s_1(t) + s_2(t)$

Considering each compartment seperately and adding the results,

$$\frac{ds}{dt} = Ax - E_1(t) = Ax - k P_1(t) = Ax - as,$$

hence  $s_1(t) = \frac{Ax}{a} (1 - e^{-at})$  and  $s_2(t) = \frac{Bx}{b} (1 - e^{-bt})$

Since  $E(t) = as_1 + Bs_2$

$$P(t) = \frac{1}{k} (as_1 + bs_2)$$

DOE ARCHIVES

equilibrium body burden  $M = \lim_{t \rightarrow \infty} s(t)$

$$t \rightarrow \infty$$

$$s(t) = x \left[ \frac{A}{a} (1 - e^{-at}) + \frac{B}{b} (1 - e^{-bt}) \right]$$

$$M = x \left( \frac{A}{a} + \frac{B}{b} \right)$$

$$P(t) = \frac{x}{k} \left[ A (1 - e^{-at}) + B (1 - e^{-bt}) \right] = \frac{x}{k} (1 - Ae^{-at} - Be^{-bt})$$

$$1. \quad y = \% \text{ of equilibrium} = \frac{s(t)}{M} = \frac{\frac{A}{a} (1 - e^{-at}) + \frac{B}{b} (1 - e^{-bt})}{\frac{A}{a} + \frac{B}{b}}$$

$$2. \quad s(t) = kP(t) \frac{\frac{A}{a} (1 - e^{-at}) + \frac{B}{b} (1 - e^{-bt})}{A(1 - e^{-at}) + B(1 - e^{-bt})}$$

$$3. \quad x = \frac{kP(t)}{A(1 - e^{-at}) + B(1 - e^{-bt})}$$

$$4. \quad M = kP(t) \frac{\frac{A}{a} + \frac{B}{b}}{A(1 - e^{-at}) + B(1 - e^{-bt})}$$

f. The following values for strontium metabolism were obtained from Supplement

No. 6 of the British Journal of Radiology:

$$A = \frac{7}{12} \quad B = \frac{5}{12} \quad k = 1 \quad \text{and} \quad s_0 = 0$$

$a = 3.65 \times 10^{-3} \text{ days}^{-1}$  and  $b = 1.73 \times 10^{-4} \text{ days}^{-1}$ , corresponding to a half-time of elimination of 190 and 4000 days respectively.

$x = 0.6 x'$  and  $x'$  is total daily intake.

At  $t = 270$  days:

$$P(t) = 3.45 \times 10^{-6} \text{ uc/day (1954 Exposed Rongelap Subjects)}$$

$$3.9 \times 10^{-6} \text{ uc/day (control Rongelap Subjects - Unexposed 1954)}$$

II. In the case of cesium and zinc:

$$s(t) = \frac{x}{a} (1 - e^{-at}) + s_0 e^{-at}$$

x is the daily accretion in uc/day, and  $s_0$  is the body burden in uc at  $t = 0$ .

$$\begin{aligned} \frac{ds}{dt} &= -as + x = -a \left[ s_0 e^{-at} + \frac{x}{a} (1 - e^{-at}) \right] + x \\ &= -E(t) + x \end{aligned}$$

$$kP(t) = E(t) = a \left[ s_0 e^{-at} + \frac{x}{a} (1 - e^{-at}) \right]$$

$$M = \lim_{t \rightarrow \infty} s(t) = \frac{x}{a}$$

Zinc<sup>65</sup>

$$a = 6.3 \times 10^{-3} \text{ days}^{-1} \quad (t_{1/2} = 110 \text{ days}) \quad k = 10$$

$$s_0 = 0.03 \text{ uc}$$

Cesium<sup>137</sup>

$$a = 4.6 \times 10^{-3} \text{ days}^{-1} \quad (t_{1/2} = 150 \text{ days}) \quad k = 1\#$$

$$s_0 = 0$$

#The urinary/fecal ratio of radiocesium for human subjects is about 5/1, so that estimates of body burden are too low by about 20 percent.